

G. Cornuéjols · C. Lemaréchal

A convex-analysis perspective on disjunctive cuts [★]

Received: September 28, 2004 / Accepted: August 31, 2005
Published online: 10 November 2005 – © Springer-Verlag 2005

Abstract. We treat with tools from convex analysis the general problem of cutting planes, separating a point from a (closed convex) set P . Crucial for this is the computation of extreme points in the so-called reverse polar set, introduced by E. Balas in 1979. In the polyhedral case, this enables the computation of cuts that define facets of P . We exhibit three (equivalent) optimization problems to compute such extreme points; one of them corresponds to selecting a specific normalization to generate cuts. We apply the above development to the case where P is (the closed convex hull of) a union, and more particularly a union of polyhedra (case of disjunctive cuts). We conclude with some considerations on the design of efficient cut generators. The paper also contains an appendix, reviewing some fundamental concepts of convex analysis.

Key words. Integer programming – Cutting planes – Separation – Disjunctive cut – Lift and project – Convex analysis

1. Introduction

An important tool in combinatorial optimization is cut generation:

- (i) a linear function $c^\top x$ must be maximized over a “hard-to-deal-with” set $S \subset \mathbb{R}^n$ (typically a huge set of integer points);
- (ii) for want of a tractable method, S is relaxed to some “easier” set R , usually polyhedral, over which $c^\top x$ is maximized;
- (iii) this may produce some \bar{x} not even belonging to $\overline{\text{conv}}(S)$, the closed convex hull of S ;
- (iv) then the problem is to separate \bar{x} from S – i.e. from $\overline{\text{conv}}(S)$ – by some *cutting plane(s)*, in order to describe S more accurately.

A cutting plane is defined by the equation $d^\top x = \kappa$, where the nonzero direction $d \in \mathbb{R}^n$ and the righthand side $\kappa \in \mathbb{R}$ satisfy

$$d^\top \bar{x} > \kappa \quad \text{and} \quad d^\top x \leq \kappa \quad \text{for all } x \in S$$

or equivalently: $d^\top \bar{x} > \sigma_S(d)$, where σ_S is the so-called *support function* of S (some essential concepts from convex analysis are given in the Appendix).

G. Cornuéjols: Tepper School of Business, Carnegie Mellon University, Pittsburgh, PA, 15213, LIF, Faculté des Sciences de Luminy, 13288 Marseille, France. e-mail: gc0v@andrew.cmu.edu

C. Lemaréchal: Inria, 655 avenue de l'Europe, Montbonnot, 38334 Saint Ismier, France. e-mail: claude.lemarechal@inrialpes.fr

Mathematics Subject Classification (2000): 52A41, 90C11

[★] Supported by NSF grant DMII-0352885, ONR grant N00014-03-1-0188, INRIA grant ODW and IBM.

Because S is hard to deal with directly, a possible approach for constructing cutting planes is to introduce an intermediate set $P \supset S$ not containing \bar{x} and to find d separating \bar{x} from P . Some desirable properties for a good cut are as follows:

- It should touch P , which fixes κ to its smallest possible value: $\kappa = \sigma_P(d)$. Thus, finding a cut is a problem in $d \in \mathbb{R}^n$ only.
- It should be as deep as possible, i.e. $d^\top \bar{x} - \sigma_P(d)$ should be maximal in a certain sense. Naturally, this requires in particular an appropriate *normalization* of d (to avoid obtaining $d^\top \bar{x} - \sigma_P(d) = +\infty$); several works are devoted to defining suitable normalizations [2, 6, 3, 11].
- When the enlargement P is polyhedral, a good cut should also touch P on a *facet*; this means that the face $F_P(d)$ of P exposed by d (see (A.4) in the Appendix) should have maximal dimension.

The present paper is thus devoted to the general problem of defining good cuts according to the above criteria; and for this we use the language and tools from convex analysis.

An application that we have in mind is in integer programming. For example in the case of pure 0-1 programming, S has the form $S := R \cap \{0, 1\}^n$ for some given polyhedron $R \subset [0, 1]^n$. Optimizing $c^\top x$ over the relaxation R yields some point \bar{x} . When $\bar{x} \in \{0, 1\}^n$, the integer program is solved. When this is not the case, then $0 < \bar{x}_j < 1$ for some j , and a natural choice for the above-mentioned enlargement P of S is the (convex hull of the) union of the polyhedra

$$R \cap \{x : x_j = 0\} \quad \text{and} \quad R \cap \{x : x_j = 1\}.$$

Note: if \bar{x} is an extreme point of R , it cannot lie in this convex hull; the inequalities that separate \bar{x} from this set P are called the *lift-and-project cuts* [2]. In practice, a small set of lift-and-project cuts with the good characteristics mentioned earlier are appended to the definition of R , thus strengthening the relaxation. The process can be repeated several times for further strengthening.

In a mixed integer linear program, $S := R \cap (\mathbb{N}^p \times \mathbb{R}^q)$ where p, q are positive integers such that $p+q = n$, and $R \subset \mathbb{R}^n$ is a polyhedron. If the point \bar{x} obtained by optimizing $c^\top x$ over R does not belong to $\text{conv}(S)$, one may strengthen the relaxation R by adding cuts, such as *split cuts* [8] defined as follows. Consider a disjunction “ $\pi^\top x \leq \pi_0$ or $\pi^\top x \geq \pi_0 + 1$ ” satisfied by all points of S but not by \bar{x} . A particular choice that gives a valid disjunction for S is $(\pi, \pi_0) \in \mathbb{N}^{n+1}$ with $\pi_j = 0$ for $j = p + 1, \dots, n$. The set P is the union of

$$R \cap \{x : \pi^\top x \leq \pi_0\} \quad \text{and} \quad R \cap \{x : \pi^\top x \geq \pi_0 + 1\}$$

and split cuts are inequalities that separate \bar{x} from P . A lift-and-project cut is a special type of split cut, where π is a unit vector. Another well-known example of split cut is Gomory’s mixed integer cut [9].

More generally, disjunctive programming, as was introduced by Balas in the 1970’s [1], considers a union of polyhedra. Specifically, given a finite number of polyhedra P_i , one would like to separate \bar{x} from $\cup_i P_i$ or equivalently from $P := \text{conv}(\cup_{i=1}^k P_i)$. We will refer to P as the *disjunctive polyhedron*. Unfortunately, a description of the disjunctive polyhedron as $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ may require an exponential number of

constraints $Ax \leq b$. Balas' key result is that P has a compact representation in a higher-dimensional space, namely there is a polyhedron $\tilde{P} := \{(x, y) \in \mathbb{R}^{n+p} : Cx + Dy \leq d\}$, whose projection onto the x -space is P , and \tilde{P} has only around kn variables and $\sum_{i=1}^k m_i$ constraints; here k is the number of polyhedra and m_i denotes the number of constraints of P_i . As a consequence, separating \bar{x} from P can be solved efficiently by linear programming. We also mention [7], in which unions of non-polyhedral sets are considered.

The paper is organized as follows. Sections 2 to 4 introduce from different points of view the fundamental objects for cutting: the *reverse polar set* and its associated support function. In §5 they are particularized to unions of sets and the polyhedral case is studied in §6. Finally we have a conclusion section: it turns out that good cuts between \bar{x} and P are attached to the direction pointing from \bar{x} to some point $x^* \in P$, and we suggest some possibilities for choosing x^* . Essential concepts from convex analysis are gathered in an Appendix.

2. Reverse polar and gauge

In this section we consider a nonempty set $Q \subset \mathbb{R}^n$ (standing for $P - \bar{x}$), which we want to separate from the origin. As observed in the introduction, this amounts to finding a direction d such that $\sigma_Q(d) < 0$. Separating directions thus form an open set, which is delicate to manage numerically, when one wants to choose a distinguished element in it. The following definition, given in [1], ingeniously turns the difficulty around, taking advantage of positive homogeneity: the set

$$Q^- := \{d \in \mathbb{R}^n : \sigma_Q(d) \leq -1\} \tag{2.1}$$

will be called the *reverse polar* of Q . Note that Q^- is dual to the space where Q lies; if “ -1 ” in (2.1) were replaced by 0 , we would recognize the definition of a polar cone; see Fig. 2.1, where the notation $\overline{\text{cone}}(\cdot)$ stands for the closed conical hull.

Theorem 2.1. *Assume $0 \notin \overline{\text{conv}}(Q)$.*

- (i) *The reverse polar Q^- is nonempty closed and convex.*
- (ii) *Let $\text{lin}(Q)$ be the linear space generated by Q . Then Q^- has the form $R + (\text{lin}(Q))^\perp$, where $R \subset \text{lin}(Q)$.*
- (iii) *A direction d separates 0 from Q if and only if $td \in Q^-$ for some $t > 0$.*




Fig. 2.1. Reverse polar (dotted lines connote dual elements)

Proof. (i) Closedness and convexity hold because the function σ_Q is itself closed and convex; see the Appendix for notation. Nonemptiness is just standard separation between $\overline{\text{con}}(Q)$ and $\{0\}$.

(ii) The property $\ell \perp \text{lin}(Q)$ is equivalent to

$$\ell^\top x = 0, \text{ i.e. } (d + \ell)^\top x = d^\top x, \text{ for all } x \in Q.$$

Thus $\sigma_Q(d) = \sigma_Q(d + \ell)$ and therefore $d \in Q^- \Leftrightarrow d + \ell \in Q^-$.

(iii) Finally, d separates 0 from Q means that $\sigma_Q(d) \leq -1/t$, for some $t > 0$; this is equivalent to $\sigma_Q(td) \leq -1$, i.e. $td \in Q^-$. □

It is observed in [2] – and will be confirmed below – that, to compute good separating directions, it is advisable to maximize some linear function over the reverse polar set. This motivates the following definition and notation:

Definition 2.2. *The support function of Q^-*

$$\mathbb{R}^n \ni q \mapsto \gamma_Q(q) := \sup_{\sigma_Q(d) \leq -1} q^\top d \tag{2.2}$$

will be called the reverse gauge of Q : $\gamma_Q = \sigma_{Q^-}$.

As explained in (A.4) of the Appendix, the (possibly empty) optimal set in (2.2) is the face of Q^- exposed by q :

$$\Gamma_Q(q) := F_{Q^-}(q) = \underset{d \in Q^-}{\text{Argmax}} q^\top d. \tag{2.3} \quad \square$$

The following result goes along the lines of [3, Thm. 12].

Theorem 2.3. *Assume $0 \notin \overline{\text{con}}(Q)$.*

- (i) *The reverse gauge is a closed sublinear function.*
- (ii) *Denoting by $\overline{\text{con}}(Q)$ the closed conical hull of Q , there holds*

$$\gamma_Q(q) \begin{cases} \leq 0 & \text{if } q \in \overline{\text{con}}(Q) \\ = +\infty & \text{otherwise.} \end{cases}$$

(iii) *If $q \in \overline{\text{con}}(Q)$ then $\gamma_Q(q) \leq -1$.*

Proof. (i) Being a support function, γ_Q is closed sublinear.

(ii) Use the notation $K := \overline{\text{con}}(Q)$ and observe that $Q^- \subset K^\circ$: if $q \in K$, then the objective value in (2.2) is nonpositive, as well as its supremum.

On the other hand, let $q \notin K$: there exists $d_0 \in K^\circ$ such that $q^\top d_0 > 0$ (separation between q and K , see (A.3) in the Appendix). Note: $\sigma_Q(d_0) \leq \sigma_K(d_0) \leq 0$ (see (A.8) in the Appendix). Take $\bar{d} \in Q^- \subset K^\circ$, $\varepsilon > 0$ and form the vector

$$d(\varepsilon) := d_0 + \varepsilon \bar{d} \in K^\circ.$$

For ε small enough, $q^\top d(\varepsilon) > 0$. By subadditivity of σ_Q , we also have

$$\sigma_Q(d(\varepsilon)) \leq \sigma_Q(d_0) + \varepsilon \sigma_Q(\bar{d}) \leq -\varepsilon.$$

Then, for $t \rightarrow +\infty$, we have

$$\sigma_Q(td(\varepsilon)) \rightarrow -\infty \quad \text{and} \quad q^\top(td(\varepsilon)) \rightarrow +\infty.$$




Fig. 2.2. Constructing a separator via a primal point

(iii) If $q \in \overline{\text{conv}}(Q)$ then $q^\top d \leq \sigma_Q(d)$ for all d (separation theorem, see (A.3) in the Appendix). Thus $q^\top d \leq -1$ for any d feasible in (2.2) and this inequality is still valid for the supremum $\gamma_Q(q)$. \square

Figure 2.2 illustrates Definition 2.2: shifting the hyperplane q^\perp so as to just touch the dashed area gives $\Gamma_Q(q)$ as the contact set. These various dual objects are all associated with a single primal point q .

We finish this section with a comment on our nonemptiness assumption. The case $Q = \emptyset$ is somewhat trivial but also pathological: we may set $\sigma_Q \equiv -\infty$ (see Note 2, p.584) and it is natural from (2.1) to set $Q^- = \mathbb{R}^n$; then $\gamma_Q(0) = 0$ and $\gamma_Q(d) = +\infty$ for all $d \neq 0$.

3. Geometric characterizations

This section is devoted to describing γ_Q and Γ_Q in terms of Q itself, without any reference to σ_Q . This can be done via the following problem:

$$\begin{cases} \sup t \\ t \geq 0 \\ q \in t \overline{\text{conv}}(Q) \end{cases} \tag{3.1}$$

and Theorem 3.2 below will justify our terminology “reverse gauge” (see the Appendix for the definition of a gauge).

In the remainder of this section, we will assume $q \in \overline{\text{conv}}(Q)$, which is the case of interest for the applications we have in mind. Observe that, by positive homogeneity, this amounts to taking q anywhere in the cone generated by Q ; up to the closure operation, this essentially amounts to assuming $\gamma_Q(q) < +\infty$ (see Theorem 2.3 (ii)).

Lemma 3.1. *Assume $0 \notin \overline{\text{conv}}(Q)$ and let $q \in \overline{\text{conv}}(Q)$. Then the optimal t in (3.1) is not smaller than 1 and the optimal value is attained (the sup is actually a max).*

Proof. Call t^* the optimal value of (3.1). The property $t^* \geq 1$ is obvious by our choice of q .

Let t_k be a maximizing sequence: $q/t_k \in \overline{\text{conv}}(Q)$ and $1/t_k \rightarrow 1/t^*$ (a finite number, possibly 0). Then $q/t_k \rightarrow q/t^*$, which has to lie in the closed set $\overline{\text{conv}}(Q)$. \square

Theorem 3.2. Assume $0 \notin \overline{\text{conv}}(Q)$ and let $q \in \overline{\text{conv}}(Q)$. Then the optimal value in (3.1) is $-\gamma_Q(q)$.

Proof. Apply Lagrangian duality to (2.2) (a convex problem which satisfies Slater’s condition):

$$\gamma_Q(q) = \inf_{t \geq 0} \sup_{d \in \mathbb{R}^n} [q^\top d - t(\sigma_Q(d) + 1)] = \inf_{t \geq 0} [-t + \theta(t)],$$

where $\theta(t) := \sup_d [q^\top d - t\sigma_Q(d)]$. Observe that $\theta(t) \geq 0$ (just take $d = 0$). Because $\gamma_Q(q) \leq -1$ (Theorem 2.3), we may impose the constraint $t \geq 1$. Then

$$\theta(t) = \begin{cases} 0 & \text{if } q \in tQ, \\ +\infty & \text{otherwise.} \end{cases}$$

To see this, either invoke [10, §E.1] (more specifically Example 1.1.5 and Theorem 1.3.5) or argue as follows: by positive homogeneity $\theta(t) = +\infty$ if there is d such that $q^\top d - t\sigma_Q(d) > 0$, i.e. $q/t \notin Q$ (separation between Q and q/t , (A.3) in the Appendix). On the other hand $q/t \in Q$ means $(q^\top/t)d - \sigma_Q(d) \leq 0$ for all d and then $\theta(t) = 0$.

It follows that $\gamma_Q(q) = \inf \{-t : t \geq 0, q \in tQ\}$. □

This result, together with Lemma 3.1, reveals important objects attached to Q and q :

Definition 3.3. Denoting by $t_Q^*(q)$ the optimal value of (3.1),

$$q^* := \frac{q}{-\gamma_Q(q)} = \frac{q}{t_Q^*(q)}$$

is the visible point from the origin in the direction q . □

Geometrically, q^* is the closest point to the origin on the line-segment $[0, q]$, still lying in $\overline{\text{conv}}(Q)$; see [3, Corr. 14]. In fact, (3.1) can also be written (setting $s := 1/t$):

$$\inf s, \quad s \geq 0, \quad sq \in \overline{\text{conv}}(Q).$$

Clearly enough, $q \mapsto t_Q^*(q)$ is positively homogeneous: the length of q has no real influence on (2.2) or (3.1) or q^* . The points $q \in \overline{\text{conv}}(Q)$ that are visible from the origin are those such that $t_Q^*(q) = -\gamma_Q(q) = 1$.

Theorem 3.4. Assume $0 \notin \overline{\text{conv}}(Q)$, let $q \in \overline{\text{conv}}(Q)$ and use the notation of Definition 3.3. Then $\Gamma_Q(q)$ is the set of d in the normal cone to $\overline{\text{conv}}(Q)$ at q^* that satisfy $q^\top d = \gamma_Q(q)$. In other words, $d \in \Gamma_Q(q)$ is equivalent to

$$d^\top(x - q^*) \leq 0 \text{ for all } x \in Q \tag{3.2}$$

$$\text{and } d^\top q = \gamma_Q(q). \tag{3.3}$$




Fig. 3.1. The visible point q^* lies in the face of Q exposed by $\Gamma_Q(q)$

Proof. Let d satisfy (3.2); maximizing over $x \in Q$ shows that $\sigma_Q(d) \leq d^\top q^*$. If, in addition, d satisfies (3.3), $\sigma_Q(d) \leq -1$: d is feasible in (2.2); use again (3.3) to conclude that d is also optimal.

Conversely, let $d \in \Gamma_Q(q)$; then $\sigma_Q(d) \leq -1$ and $q^\top d = \gamma_Q(q)$, hence

$$\sigma_Q(d) \leq -1 = d^\top q^* . \tag{3.4}$$

However, $q^* \in \overline{\text{conv}}(Q)$ (Lemma 3.1 and Theorem 3.2). By definition of a support function, $\sigma_Q(d) \geq d^\top q^*$: equality holds in (3.4), i.e. q^* lies in the face of Q exposed by d (see (A.4) in the Appendix);

$$q^* \in F_{\overline{\text{conv}}(Q)}(d) , \quad \text{i.e.} \quad d \in N_{\overline{\text{conv}}(Q)}(q^*) ,$$

where the last equivalence is (A.6) in the Appendix. □

Thus, a direction $d \in \Gamma_Q(q)$ exposes q^* in $\overline{\text{conv}}(Q)$. Said differently, $d \in \Gamma_Q(q)$ defines a hyperplane

$$\{x \in \mathbb{R}^n : d^\top x = \sigma_Q(d)\}$$

supporting $\overline{\text{conv}}(Q)$ at q^* . The important property to characterize $\Gamma_Q(q)$ is (3.2), which expresses that it lies in the normal cone to Q at q^* (the thick dashed arrow of Fig. 3.1); while (3.3) just acts as a normalization (specifying the length of the arrow).

The constructions given in this section are intrinsic: they do not rely on any representation of Q . It can be mentioned that the dilation effect revealed in (3.1) may result in rather intricate calculations when Q is represented by nonlinear constraints – see [7]. For affine constraints, the calculations considerably simplify – see §6 below, more specifically Remark 6.5.

4. Normalizations

The reverse gauge can also be computed via a convex optimization problem with a single linear constraint, namely

$$\inf \sigma_Q(d) , \quad q^\top d \geq -1 ; \tag{4.1}$$

observe that this problem would become very similar to (2.2) if the righthand side “−1” were replaced by $\gamma_Q(q)$. In fact, the aim of the present section is to show that the optimal value κ^* and optimal set D^* in (4.1) are just proportional to the optimal value γ and optimal set Γ in Definition 2.2. The profound reason is positive homogeneity: replacing “−1” by any $\kappa < 0$ in the definition (2.1) of the reverse polar would change essentially nothing. As a result, changing the righthand side in (2.2) or (4.1) just amounts to dilating the picture. On Fig. 2.2, one could also shift q^\perp by a fixed amount, $\Gamma_Q(q)$ could be recovered by an appropriate dilation of Q^- .

To prove equivalence between (2.2) and (4.1), we first show that their constraint can be turned to an equality:

Lemma 4.1. *Assume $0 \notin \overline{\text{conv}}(Q)$ and let $q \in \overline{\text{conv}}(Q)$. Then:*

(i) $\gamma_Q(q)$ and $\Gamma_Q(q)$ are the optimal value and optimal set of

$$\max q^\top d, \quad \sigma_Q(d) = -1;$$

(ii) $\kappa^* < 0$ and D^* are the optimal value and optimal set of

$$\min \sigma_Q(d), \quad q^\top d = -1.$$

Proof. Let d be feasible in (2.2): $\sigma_Q(d) =: \alpha \leq -1$; by definition of a support function, $q^\top d \leq \sigma_Q(d) < 0$. By positive homogeneity, $\sigma_Q(\frac{d}{-\alpha}) = -1$ and $q^\top \frac{d}{-\alpha} \geq q^\top d$, with strict inequality if $-\alpha > 1$. This proves (i). The proof of (ii) goes the same way. \square

This enables us to make the equivalence precise:

Theorem 4.2. *Assume $0 \notin \overline{\text{conv}}(Q)$ and let $q \in \overline{\text{conv}}(Q)$. Then*

$$\gamma_Q(q) = \frac{1}{\kappa^*} \quad \text{and} \quad \Gamma_Q(q) = \frac{D^*}{-\kappa^*}.$$

Proof. We use the shortened notation $\gamma := \gamma_Q(q)$. Recall that $\gamma \leq -1$ (Theorem 2.3(iii)).

Let $\varepsilon > 0$. In view of Lemma 4.1(i), we can take d_ε such that $\sigma_Q(d_\varepsilon) = -1$ and $q^\top d_\varepsilon \geq \gamma - \varepsilon$. Then set $d := \frac{d_\varepsilon}{-\gamma + \varepsilon}$, so that $q^\top d \geq -1$: d is feasible in (4.1), hence

$$\kappa^* \leq \sigma_Q(d) = \frac{1}{-\gamma + \varepsilon} \sigma_Q(d_\varepsilon) = \frac{1}{\gamma - \varepsilon}.$$

Since $\varepsilon > 0$ is arbitrary, $\kappa^* \leq 1/\gamma$, i.e. $\kappa^* \gamma \geq 1$, i.e. $\gamma \leq 1/\kappa^*$.

On the other hand, Lemma 4.1(ii) allows us to force $\varepsilon \in]0, -\kappa^*[$ and to take d'_ε such that $q^\top d'_\varepsilon = -1$ and $\sigma_Q(d'_\varepsilon) \leq \kappa^* + \varepsilon < 0$. Then set $d := \frac{d'_\varepsilon}{-\kappa^* - \varepsilon}$, so that $\sigma_Q(d) \leq -1$: d is feasible in (2.2), hence

$$\gamma \geq q^\top d = q^\top \frac{d'_\varepsilon}{-\kappa^* - \varepsilon} = \frac{1}{\kappa^* + \varepsilon}.$$

Since $\varepsilon > 0$ is arbitrarily small, $\gamma \geq 1/\kappa^*$. Altogether, $\gamma = 1/\kappa^*$.

Now $d \in \Gamma_Q(q)$ means

$$q^\top d = \gamma \quad \text{and} \quad \sigma_Q(d) = -1,$$

while $d \in D^*$ means

$$q^\top d = -1 \quad \text{and} \quad \sigma_Q(d) = \frac{1}{\gamma}.$$

The relation between $\Gamma_Q(q)$ and D^* follows by positive homogeneity. □

With reference to our introduction, the constraint in (4.1) can be viewed as a normalization: to construct a tightest cut d amounts to minimizing σ_Q and requires a bound on d . The above result establishes that bounding $q^\top d$ (from below) does the job: it imposes the simplest possible constraint to give a meaning to the minimization of σ_Q . It does not necessarily produce a separating direction, though: consider in \mathbb{R}^2 the circle

$$Q := \{x = (\xi, \eta) : (\xi - 1)^2 + (\eta - 1)^2 \leq 1\}$$

and take $q = (1, 0)$. Then $\gamma_Q(q) = -1$ (Theorem 3.2). However, no $d \in (0, \mathbb{R}_-) = N_Q(q)$ satisfies $d^\top q = -1$: in fact $\Gamma_Q(q) = \emptyset$.

Remark 4.3. The proof of Theorem 3.2 shows that (3.1) is the Lagrangian dual of (2.2). A similar interpretation applies to (4.1) through the following formulation of (3.1):

$$\inf_{x \in Q, t \geq 0} t, \quad tq = x.$$

Dualizing the constraint $tq - x = 0$ with multiplier d , we obtain the Lagrangian dual

$$\sup_{d \in \mathbb{R}^n} \inf_{x \in Q, t \geq 0} t + d^\top (tq - x).$$

The internal inf-problem can be solved separately in x and t . Remembering the definition of σ_Q , we obtain

$$\sup_{d \in \mathbb{R}^n} -\sigma_Q(d), \quad q^\top d + 1 \geq 0,$$

which is nothing other than (4.1).

Note also the optimality condition of

$$\inf_{x \in Q} -d^\top x,$$

which is $d \in N_Q(x)$ and “explains” Theorem 3.4. □




Fig. 4.1. No separator exposes q^*

5. The case of a disjunction

Let Q_0 and Q_1 be two nonempty subsets of \mathbb{R}^n . In this section, we take $Q = Q_0 \cup Q_1$ and we apply the above results to compute the reverse elements of Q in terms of the data Q_0 and Q_1 .

Lemma 5.1. *Denote by σ_i and γ_i the support function and reverse gauge of Q_i , $i = 0, 1$. There holds*

$$\sigma_Q = \max \{ \sigma_0, \sigma_1 \}, \quad Q^- = Q_0^- \cap Q_1^-, \quad \gamma_Q = \overline{\text{conv}} (\min \{ \gamma_0, \gamma_1 \}).$$

Proof. See the Appendix for σ_Q . The formula for Q^- follows easily. See again the Appendix for $\gamma_Q = \sigma_{Q^-}$. □

In the following result, the notation \bar{S} is used for the closure of a set S .

Theorem 5.2. *Assume $0 \notin \overline{\text{conv}} (Q)$ and let $q \in \overline{\text{conv}} (Q)$. Then $-\gamma_Q(q)$ is the optimal value of*

$$\begin{cases} \sup t_0 + t_1 \\ t_0 \geq 0, t_1 \geq 0 \\ q \in \overline{t_0 \text{conv} (Q_0) + t_1 \text{conv} (Q_1)}. \end{cases}$$

Proof. Apply Theorem 3.2: $-\gamma_Q(q)$ is the optimal value of (3.1). Now $t \overline{\text{conv}} (Q)$ is clearly $\overline{\text{conv}} (tQ)$, which is the closure of $S_t := \text{conv} (tQ)$. Using associativity of convex combinations, we can write

$$S_t = \{ \alpha t q_0 + (1 - \alpha) t q_1 : \alpha \in [0, 1], q_0 \in \text{conv} (Q_0), q_1 \in \text{conv} (Q_1) \}.$$

Introduce the notation $t_0 := \alpha t$, $t_1 := (1 - \alpha) t$. When α describes the segment $[0, 1]$, t_0 and t_1 describe the corresponding segment dilated by $t \geq 0$, so that

$$S_t = \{ t_0 \text{conv} (Q_0) + t_1 \text{conv} (Q_1) : t_0 \geq 0, t_1 \geq 0, t_0 + t_1 = t \}. \quad \square$$

Introducing the visible point q^* of Definition 3.3 allows the characterization of $\Gamma_Q(q)$ in terms of normal cones (see (A.5) in the Appendix):

Theorem 5.3. *Assume $0 \notin \overline{\text{conv}} (Q)$ and let $q \in \overline{\text{conv}} (Q)$. Then $\Gamma_Q(q)$ is the set of d satisfying*

$$\begin{aligned} d^\top (x - q^*) &\leq 0 \text{ for all } x \in Q_0, \\ d^\top (x - q^*) &\leq 0 \text{ for all } x \in Q_1, \\ \text{and } q^\top d &= \gamma_Q(q). \end{aligned}$$

Proof. Use Theorem 3.4 and observe that, if d satisfies $d^\top (x - q^*) \leq 0$ for all $x \in Q_i$, $i = 0, 1$, then the same inequality holds

- for all $x \in \text{conv} (Q_0 \cup Q_1)$ by convex combinations,
- and for all x in the closure of the latter set, by continuity.

Altogether, (3.2) is satisfied. The converse is obvious. □




Fig. 5.1. Apexed convex hull and corresponding normal cone

Remark 5.4. This result suggests that the normal cone $N_Q(q^*)$ (where Q is a union) is the intersection of two appropriate normal cones, as illustrated by Fig. 5.1: for $i = 1, 2$, we have

$$Q_i^* := \text{conv}(Q_i \cup \{q^*\}) = \{(1 - \alpha)q^* + \alpha x : x \in Q_i, \alpha \in [0, 1]\} \\ = q^* + \{\alpha(Q_i - q^*) : \alpha \in [0, 1]\}.$$

It follows that the directions d satisfying

$$d^\top(x - q^*) \leq 0 \text{ for all } x \in Q_i,$$

or equivalently

$$d^\top \alpha(x - q^*) \leq 0 \text{ for all } x \in Q_i \text{ and } \alpha \in [0, 1]$$

just make up the normal cone $N_{Q_i^*}(q^*)$ to Q_i^* at q^* . As a result, $d \in \Gamma_Q(q)$ means

$$d \in N_{Q_0^*}(q^*) \cap N_{Q_1^*}(q^*) \\ \text{and } q^\top d = \gamma_Q(d).$$

The whole issue when dealing with a disjunction is thus to characterize a normal cone of the type $N_{\text{conv}(S \cup \{0\})}(0)$, for some set S (here $S = Q_i - q^*$). Such a cone can also be viewed as the polar of cone (S) . □

One can also compute the reverse elements of a disjunction using a normalization, as in §4:

Theorem 5.5. Denote by κ^* and D^* the optimal value and optimal set (possibly empty) of

$$\inf \kappa, \quad \kappa \geq \sigma_0(d), \quad \kappa \geq \sigma_1(d), \quad q^\top d = -1.$$

Then $\gamma_Q(q) = \frac{1}{\kappa^*}$ and $\Gamma_Q(q) = \frac{D^*}{-\kappa^*}$.

Proof. Use Lemma 5.1 in (4.1): the result is a straightforward application of Theorem 4.2. □

6. Polyhedra

A polyhedron can be described in two different ways: as a convex hull (Theorem 6.1 below), or as an intersection of half-spaces, as in §6.1. The case of a disjunction (§6.2) is somewhat intermediate.

Theorem 6.1. *Let Q be a nonempty closed convex polyhedron not containing the origin. Then Q^- is also a nonempty closed convex polyhedron; in particular, if $Q = \text{conv}(x^1, \dots, x^k) + \text{cone}(r^1, \dots, r^\ell)$, then*

$$d \in Q^- \iff \begin{cases} d^\top x^i \leq -1 & \text{for } i = 1, \dots, k, \\ d^\top r^j \leq 0 & \text{for } j = 1, \dots, \ell. \end{cases}$$

Furthermore, γ_Q is a piecewise linear function in the domain where it is finite; $\Gamma_Q(q)$ is a closed convex polyhedron, nonempty if $q \in \overline{\text{cone}}(Q) = \text{cone}(Q)$.

Proof. Everything is rather clear; in particular, (2.2) is a linear program, which has a nonempty optimal set if it has a finite optimal value. □

Now comes a result motivating this theory.

Theorem 6.2. *The face $F_Q(d)$ exposed by $d \in Q^-$ is a facet of Q if and only if*

$$d^* := \frac{d}{-\sigma_Q(d)}$$

is an extreme point of $Q^- \cap \text{lin}(Q)$.

Proof. The points $x \in Q$ that lie in $F_Q(d)$ are those such that $d^\top x = \sigma_Q(d)$, i.e. $(d^*)^\top x = -1$.

Let p denote the dimension of $\text{lin}(Q)^\perp$. Suppose $F_Q(d)$ is a facet of Q and let x^1, \dots, x^{n-p} be $n - p$ affinely independent points in $F_Q(d)$. Then d^* satisfies a system of $n - p$ independent equations $(d^*)^\top x^i = -1$ which uniquely determines it up to the linear space $\text{lin}(Q)^\perp$. Conversely, if d^* is an extreme point of $Q^- \cap \text{lin}(Q)$, there exist $n - p$ affinely independent points x^1, \dots, x^{n-p} in Q that satisfy $(d^*)^\top x^i = -1$. These points lie in $F_Q(d)$, therefore $F_Q(d)$ is a facet of Q . □

6.1. Explicitly described polyhedra

Assume that Q is described by a set of constraints:

$$Q = \{x \in \mathbb{R}^n : Ax \leq b \in \mathbb{R}^m\}. \tag{6.1}$$

A simple way of computing γ_Q is then as follows:

Theorem 6.3. *Assume Q of (6.1) does not contain the origin and let $q \in Q$; denote by A_j the j^{th} row of A . Then*

$$-\gamma_Q(q) = \max_{b_j < 0} \frac{A_j q}{b_j}.$$

Calling $J(q)$ the set of j realizing the above max, $\Gamma_Q(q)$ is the set of directions in $\text{cone}(A_j)_{j \in J(q)}$, normalized by the constraint $d^\top q = \gamma_Q(q)$.

Proof. Just apply Theorems 3.2 and 3.4: (3.1) takes the form

$$\max t, \quad t \geq 0, \quad Aq \leq tb.$$

Then use the known expression for the normal cone to a polyhedron. □

6.2. Union of polyhedra

Now take the situation of §5: $Q := \overline{\text{conv}}(Q_0 \cup Q_1)$, where

$$Q_i := \{x \in \mathbb{R}^n : A_i x \leq b_i \in \mathbb{R}^{m_i}\} \quad \text{for } i = 0, 1 \tag{6.2}$$

are nonempty closed convex polyhedra. Of course, all the constructions of the present section can be obtained via appropriate linear programs. However, we will keep close to our convex-analysis point of view.

We start by a characterization of the *closed* convex hull Q described above.

Lemma 6.4. *With the notation (6.2), $x \in Q$ if and only if there are $y_0, y_1 \in \mathbb{R}^n$ and $\alpha_0, \alpha_1 \in \mathbb{R}$ such that*

$$x = y_0 + y_1, \quad \alpha_0 + \alpha_1 = 1, \quad A_i y_i - \alpha_i b_i \leq 0 \text{ and } \alpha_i \geq 0 \text{ for } i = 0, 1. \tag{6.3}$$

Proof. Consider the set

$$S := \{\alpha_0 x_0 + \alpha_1 x_1 : \alpha_0 + \alpha_1 = 1, A_i x_i \leq b_i \text{ and } \alpha_i > 0 \text{ for } i = 0, 1\}.$$

Clearly enough, $S \subset \text{conv}(Q_0 \cup Q_1)$ and any point in $\text{conv}(Q_0 \cup Q_1)$ can be approximated to any accuracy by a point in S . It follows that Q is the closure of S .

Now set $y_i := \alpha_i x_i$ for $i = 0, 1$, so that we have

$$S = \{y_0 + y_1 : \alpha_0 + \alpha_1 = 1, A_i y_i \leq \alpha_i b_i \text{ and } \alpha_i > 0 \text{ for } i = 0, 1\},$$

whose closure is clearly the set described by (6.3). □

This result will be used to characterize tQ in §3 by appropriate constraints.

Remark 6.5. The above proof makes more precise our comments at the end of §3. In fact, suppose $Q = \{x : c(x) \leq 0\}$ (with c convex but not necessarily affine). Then the feasible set in (3.1) is given by $c(q/t) \leq 0$, a nasty nonconvex constraint (with respect to the variable (q, t)). Here comes a trick: with $t > 0$ we can also write $tc(q/t) \leq 0$ and this reveals the so-called *perspective function* of c (see [10]):

$$\mathbb{R}^n \times]0, +\infty[\ni (q, t) \mapsto \tilde{c}(q, t) := tc\left(\frac{q}{t}\right),$$

which happens to be convex (and even sublinear, see the Appendix). When c is affine, \tilde{c} is linear, which results in the simple formula (6.3). Otherwise, \tilde{c} is rather involved, which explains the technicalities developed in [7]. □

Theorem 6.6. *With the notation (6.2), assume $0 \notin Q$ and let $q \in Q$. Then $-\gamma_Q(q)$ is the optimal value of*

$$\begin{cases} \max t_0 + t_1 \\ q_0 + q_1 = q \\ A_i q_i - t_i b_i \leq 0 \text{ for } i = 0, 1 \\ t_i \geq 0 \quad \text{for } i = 0, 1. \end{cases} \tag{6.4}$$

Proof. Just dilate the set (6.3) by $t \geq 0$, replacing α_i by $t_i := t\alpha_i$. □

It remains to characterize $\Gamma_Q(q)$, i.e. the normal cone to Q at q^* . The following result, inspired by Remark 5.4, will allow the application of Theorem 5.3.

Proposition 6.7. *With the notation (6.2) and q^* of Definition 3.3, the cone*

$$N_0 := \{d \in \mathbb{R}^n : d^\top(x - q^*) \leq 0 \text{ for all } x \in Q_0\}$$

is the set of $d = A_0^\top u_0$, where $u_0 \geq 0 \in \mathbb{R}^{m_0}$ satisfies $(A_0^\top q^ - b_0)^\top u_0 \geq 0$.*

Proof. Using an artificial multiplication by $\alpha > 0$, we can write that $d \in N_0$ if and only if

$$\alpha d^\top(x - q^*) \leq 0 \quad \text{for all } x \text{ such that } A_0 x \leq b_0 \text{ and all } \alpha \geq 0$$

which, setting $y := \alpha x - \alpha q^*$, can also be written

$$d^\top y \leq 0 \quad \text{for all } \begin{pmatrix} y \\ \alpha \end{pmatrix} \in K_0;$$

here $K_0 \subset \mathbb{R}^{n+1}$ is the cone defined by $\begin{pmatrix} A_0 & A_0 q^* - b_0 \\ 0^\top & -1 \end{pmatrix} \begin{pmatrix} y \\ \alpha \end{pmatrix} \leq 0$. This means that

$$\begin{pmatrix} d \\ 0 \end{pmatrix} \in K_0^\circ.$$

Now K_0° is classically the set described by

$$\begin{pmatrix} d \\ r \end{pmatrix} = \begin{pmatrix} A_0^\top & 0 \\ (A_0 q^* - b_0)^\top & -1 \end{pmatrix} \begin{pmatrix} u_0 \\ v_0 \end{pmatrix} \quad \text{with } \begin{pmatrix} u_0 \\ v_0 \end{pmatrix} \geq 0 \in \mathbb{R}^{n+1}.$$

Fixing $r = 0$ introduces the constraint

$$(A_0 q^* - b_0)^\top u_0 - v_0 = 0, \quad \text{i.e. } (A_0 q^* - b_0)^\top u_0 \geq 0. \quad \square$$

The characterization of $\Gamma_Q(q)$ is now easy:

Theorem 6.8. *With the notation (6.2), assume $0 \notin Q$ and let $q \in Q$; introduce $\gamma_Q(q)$ of (6.4) and q^* of Definition 3.3. Up to the normalization $d^\top q^* = \gamma_Q(q)$, $\Gamma_Q(q)$ is the set of $d = A_0^\top u_0 = A_1^\top u_1$, with $(u_0, u_1) \in \mathbb{R}_+^{m_0} \times \mathbb{R}_+^{m_1}$ satisfying the set of constraints*

$$A_0^\top u_0 = A_1^\top u_1 \quad \text{and} \quad (A_i q^* - b_i)^\top u_i \geq 0, \quad i = 0, 1.$$

Proof. Straightforward from Theorem 5.3 and Proposition 6.7. □

Remark 6.9. It has been observed in Remark 4.3 that $\Gamma_Q(q)$ could also be obtained by applying duality to (3.1), which is here (6.4). Indeed, multiply the objective function of (3.1) by some normalizing coefficient $\kappa > 0$ and call d the multiplier associated with $q_0 + q_1 = q$. This gives the dual

$$\begin{cases} \min -q^\top d \\ d = A_i^\top u_i & \text{for } i = 0, 1 \\ \kappa + b_i^\top u_i \leq 0 & \text{for } i = 0, 1 \\ u \geq 0. \end{cases}$$

Its solution depends multiplicatively on κ , which can be adjusted so as to reproduce $q^\top d = \gamma_Q(q)$. The “correct” value is a posteriori $\kappa = -(A_0 q^*)^\top u_0 = -(A_1 q^*)^\top u_1$. \square

7. Conclusion: toward efficient cut generation

This paper has studied the construction of cutting planes defined by $(d, \sigma_Q(d)) \in \mathbb{R}^n \times \mathbb{R}$, separating the origin from (the closed convex hull of) a set Q . With regard to our introduction, this is also the problem of separating a point \bar{x} from (the closed convex hull of) a set P : it suffices to set $Q := P - \bar{x}$, and to use the obvious relation $\sigma_Q(d) = \sigma_P(d) - d^\top \bar{x}$.

The d -part of a cut is conveniently computed with the help of some $q \in \text{conv}(Q)$, or rather its corresponding visible point q^* . We solve (2.2) or (4.1) which, in the translated context, take the form

$$\sup_{\sigma_P(d) - d^\top \bar{x} \leq -1} d^\top (\underline{x} - \bar{x}) \quad \text{or} \quad \inf_{d^\top (\underline{x} - \bar{x}) \geq -1} \sigma_P(d) - d^\top \bar{x}$$

(here $\underline{x} - \bar{x}$ stands for q , with $\underline{x} \in P$).

We have not considered the problem of choosing q in Q – i.e. \underline{x} in P ; but some suggestions can be made. A guide for this choice might be the depth of a cut, defined as follows.

Definition 7.1. *The depth of a cut separating \bar{x} from a closed convex set P is the Euclidean distance between \bar{x} and the separating hyperplane. If the cut is defined by*

$$d^\top x = \sigma_{P-\bar{x}}(d), \quad \text{where } \sigma_{P-\bar{x}}(d) = \sigma_P(d) - d^\top \bar{x} < 0,$$

its depth is $\frac{-\sigma_{P-\bar{x}}(d)}{\|d\|}$. \square

In fact, P. Bonami [5] performed experiments in the context of lift-and-project cuts to solve $\max c^\top x$ over $S = \{x \in \{0, 1\}^n : Ax \leq b\}$. He compared two choices of $q = \underline{x} - \bar{x}$ by choosing $\underline{x} \in S$ in two different ways: in the first, \underline{x} was chosen to be a good heuristic solution to the integer program and in the second \underline{x} was a bad solution (obtained by minimizing $c^\top x$ instead of maximizing it). One would expect the direction q to be more central in cone (Q) in the second case. On 15 instances of Miplib 3.0¹, the average depth of a cut generated by the first choice of \underline{x} was 0.026 whereas it was 0.084

¹ <http://www.caam.rice.edu/~bixby/miplib/miplib3.html>

with the second choice. Adding cuts for all 0-1 variables x_j where \bar{x}_j was fractional gave an average improvement in objective value of 6% (of the gap between the linear and integer program objective values) in the first case and 6.6% in the second case. When the cuts were added to $Ax \leq b$ and this strengthened linear program was solved again to compute a new solution \bar{x} , this process being repeated iteratively for one minute of CPU, the difference was even more significant (38% versus 53%).

This small experiment suggests that it seems preferable to choose deep cuts, and one may therefore wish to find the deepest one; this corresponds to projecting \bar{x} onto P :

Proposition 7.2. *With the above notation, call x^* the solution of*

$$\min_{x \in P} \|x - \bar{x}\|.$$

Then $d^* := \bar{x} - x^*$ defines the deepest cut between \bar{x} and P .

Proof. Using positive homogeneity, the d -part of a cut can be restricted to a normalized d . Thus, a deepest cut is a solution of

$$\min_{\|d\|=1} \sigma_{P-\bar{x}}(d)$$

where, because the optimal objective value is negative, the constraint $\|d\| = 1$ can be replaced by $\|d\| \leq 1$. Thus, a deepest cut is a solution of

$$\min_{\|d\| \leq 1} \sup_{x \in P} d^\top (x - \bar{x}).$$

This is the min-maximization of a bilinear function over a product of closed convex sets, one of which (the unit ball) is compact. Using a standard minimax theorem, for example [12, Corollary 37.3.2], we can equivalently solve

$$\sup_{x \in P} \min_{\|d\| \leq 1} d^\top (x - \bar{x}).$$

For given $x \in P$ (and hence different from \bar{x}), the internal minimization has the unique solution

$$d(x) := \operatorname{Argmin}_{\|d\| \leq 1} d^\top (x - \bar{x}) = -\frac{x - \bar{x}}{\|x - \bar{x}\|}$$

and our minimax problem boils down to

$$\sup_{x \in P} d(x)^\top (x - \bar{x}) = \sup_{x \in P} -\|x - \bar{x}\|,$$

whose unique solution is clearly the projection x^* of \bar{x} onto P . □

Thus, to construct the deepest cut between a closed convex set P and a point $\bar{x} \notin P$, one needs to project \bar{x} onto P . When P is a polyhedron, this amounts to solving a linear-quadratic problem.

When solving an integer program by a cutting plane approach, generating just one cut is usually not enough, even the deepest one. A better strategy is to generate many cuts at once; and as already mentioned, these should preferably be facets of P . An attractive approach is therefore to operate as follows:

- (i) Compute the projection x^* of \bar{x} onto P . Set $d^* = \bar{x} - x^*$.
- (ii) Construct cuts, say

$$(d_k, \sigma_P(d_k) - \bar{x}^\top d_k), \quad k = 1, \dots, K$$

corresponding to facets of P passing through x^* .

- (iii) Append to the above list $(d^*, \sigma_P(d^*) - \bar{x}^\top d^*)$ (this step is useless if (ii) has produced enough cuts, so that d^* is a positive combination of the d_k 's).

Let us mention that Bonami [4] used the following idea for generating facets of P that define the optimum of $\max_{x \in P} c^\top x$. First find \bar{x} solving the relaxation $\max_{x \in R} c^\top x$ (we use the notation of the introduction). Then solve the linear program based on (6.4) to find a cut $(d_1, \sigma_P(d_1) - \bar{x}^\top d_1)$ separating \bar{x} from P (any $q \in Q$ will do in (6.4): for example $q = x^* - \bar{x}$ where, as above, x^* is the projection of \bar{x} onto P). Let R^1 be the relaxation obtained by adding this cut to R , and let x^1 maximize $c^\top x$ over R^1 . The procedure is then repeated: the k th iteration generates

- a cut $(d_k, \sigma_P(d_k) - \bar{x}^\top d_k)$ separating x_{k-1} from P ,
- an improved relaxation R^k obtained by adding this cut to R^{k-1} ,
- and a point x^k maximizing $c^\top x$ over R^{k-1} .

The procedure terminates when $x^k \in P$. Bonami tried this procedure on the Miplib problems mentioned earlier, for the lift-and-project relaxation

$$P_{L\&P} := \bigcap_{j=1}^n \text{conv}((R \cap \{x_j = 0\}) \cup (R \cap \{x_j = 1\})).$$

On average, maximizing $c^\top x$ over $P_{L\&P}$ closed 30% of the gap between R and $\text{conv}(S)$.

Acknowledgements. We are grateful to anonymous referees for their careful reading, which helped improve an earlier version of this paper.

A. Appendix: basic concepts from convex analysis

The convex-analysis aspects necessary for the understanding of this paper can be found in any book on modern convex analysis. We summarize here some material mainly extracted from [10, Chap. C], co-authored by the second author.

Support functions. An important object associated to a nonempty subset S of \mathbb{R}^n is its so-called support function

$$\mathbb{R}^n \ni d \mapsto \sigma_S(d) := \sup_{x \in S} d^\top x. \tag{A.1}$$

In other words, σ_S is the result of optimizing a linear function over S .

A support function enjoys the following important properties:

- (i) It is convex (being a supremum of linear functions²).
- (ii) It is positively homogeneous: $\sigma_S(td) = t\sigma_S(d)$ for all $t > 0$, so that d in (A.1) can be understood as a direction (knowing that $\sigma_S(0) = 0$).

Functions that are convex and positively homogeneous are called *sublinear*; they are in particular subadditive, i.e.

$$\sigma_S(d_1 + d_2) \leq \sigma_S(d_1) + \sigma_S(d_2).$$

Note also that σ_S is linear if and only if S is a single point. Being convex, σ_S would be continuous if it were finite everywhere. However, we may have $\sigma_S(d) = +\infty$ (when S is unbounded; for example $\sigma_{\mathbb{R}^n}(d)$ is 0 if $d = 0$ and $+\infty$ otherwise).

- (iii) Nevertheless, a support function is lower-semicontinuous, or *closed*³.

The set of functions that are sublinear and closed is important, as it directly generalizes the set of linear functions.

Correspondence between analysis and geometry. Let $\overline{\text{conv}}(S)$ denote the closed convex hull of S , equivalently defined as

- the topological closure of the convex hull of S (the closure operation is superfluous if S is already closed and bounded)
- or the intersection of all the half-spaces containing S .

Recall that a half-space associated to $(d, \beta) \in \mathbb{R}^n \times \mathbb{R}$ is defined as

$$\{x \in \mathbb{R}^n : d^\top x \leq \beta\}.$$

It is clear enough that $\sigma_S = \sigma_{\overline{\text{conv}}(S)}$, and this is the starting remark for a fundamental result: the closed convex sets S in \mathbb{R}^n are in one-to-one correspondence with the functions σ from \mathbb{R}^n to $\mathbb{R} \cup \{+\infty\}$ that are closed and sublinear. The correspondence $S \mapsto \sigma$ is given by (A.1); its inverse is given by

$$\sigma \mapsto S := \{x \in \mathbb{R}^n : \sigma(d) \geq x^\top d \text{ for all } d \in \mathbb{R}^n\} = \partial\sigma(0), \tag{A.2}$$

where the last equality is obvious, if one remembers the definition of a subdifferential:

$$\partial\sigma(\bar{d}) := \{x \in \mathbb{R}^n : \sigma(d) \geq \sigma(\bar{d}) + x^\top (d - \bar{d}) \text{ for all } d \in \mathbb{R}^n\}.$$

From the definition (A.1) itself, $d^\top x \leq \sigma_S(d)$ for all $d \in \mathbb{R}^n$ whenever $x \in \overline{\text{conv}}(S)$. The above result states that the converse is true:

$$x \in \overline{\text{conv}}(S) \iff \forall d \in \mathbb{R}^n, d^\top x \leq \sigma_S(d), \tag{A.3}$$

which is nothing other than the familiar separation theorem

$$x \notin \overline{\text{conv}}(S) \iff \exists d \in \mathbb{R}^n : d^\top x > \sigma_S(d).$$

² The assumption $S \neq \emptyset$ is convenient. We may also define $\sigma_\emptyset \equiv -\infty$, although this is a somewhat pathological “function”.

³ This means:

- its epigraph $\{(x, r) \in \mathbb{R}^n \times \mathbb{R} : r \geq \sigma_S(x)\}$ is a closed set in \mathbb{R}^{n+1} ,
- or equivalently: its sublevel sets $\{d : \sigma_S(d) \leq \beta\}$ are closed sets (possibly empty) in \mathbb{R}^n , for any β ,
- or equivalently: given arbitrary $d \in \mathbb{R}^n$ and $d_k \rightarrow d$, if $\sigma_S(d_k) \rightarrow \ell$, then $\ell \geq \sigma_S(d)$ (note: this last inequality must hold even if $\sigma_S(d) = +\infty$).

Isomorphism aspects. Not only closed convex sets and closed sublinear functions, but also *operations* between them are in correspondence via (A.1) and (A.2), which is indeed an isomorphism. For example,

$$\sigma_{S_1+S_2} = \sigma_{S_1} + \sigma_{S_2} .$$

Two other important operations are as follows: given an index set J (possibly infinite), consider closed convex sets $\{S_j : j \in J\}$ and closed sublinear functions $\{\sigma_j : j \in J\}$ related by

$$\begin{cases} \sigma_j \text{ is the support function of } S_j, \\ \text{or equivalently: } S_j = \partial\sigma_j(0) . \end{cases}$$

Form the following two pairs (S, σ) :

- (i) Set $S := \overline{\text{conv}}(\cup_j S_j)$ and $\sigma := \sup_j \sigma_{S_j}$.
- (ii) Set $S := \cap_j S_j$ and let $\sigma [:= \overline{\text{conv}}(\inf_j \sigma_j)]$ be the closed convex hull of the σ_j 's: this is the function whose epigraph is the closed convex hull of the union of the epigraphs of the σ_j 's⁴

Then, in both cases (i) and (ii) we have

$$\begin{cases} \sigma \text{ is the support function of } S, \\ \text{or equivalently: } S = \partial\sigma(0) . \end{cases}$$

Exposed faces and normal cones. Here S is assumed to be closed and convex. A half-space $\{x : d^\top x \leq \beta\}$ contains S if and only if $\beta \geq \sigma_S(d)$. If $\beta = \sigma_S(d)$ and if the supremum in (A.1) is attained, then the hyperplane $\{x : d^\top x = \sigma_S(d)\}$ is said to support S . The contact set

$$F_S(d) := \underset{x \in S}{\text{Argmax}} d^\top x = \{x \in S : d^\top x = \sigma_S(d)\} \tag{A.4}$$

is called the *face* (of S) *exposed* by d (note: $F_S(0) = S$).

Knowing that the *normal cone* to S at $x \in S$ is

$$N_S(x) := \{d \in \mathbb{R}^n : d^\top (x' - x) \leq 0 \text{ for all } x' \in S\} , \tag{A.5}$$

the following equivalence is clear:

$$x \in F_S(d) \iff d \in N_S(x) . \tag{A.6}$$

⁴ The epigraph of σ is thus the intersection of the half-spaces containing the epigraphs of all the σ_j 's: the property $r \geq \sigma(d)$ means

$$\begin{aligned} r &\geq \alpha^\top d + \beta && \text{for all } \alpha, \beta \text{ such that} \\ &&& \alpha^\top d' + \beta \leq \sigma_j(d') \text{ for all } d' \in \mathbb{R}^n \text{ and all } j \in J . \end{aligned}$$

Gauges and polar sets. The polar of a set $S \subset \mathbb{R}^n$ is

$$S^\circ := \{y : y^\top x \leq 1 \text{ for all } x \in S\} = \{y : \sigma_S(y) \leq 1\}. \quad (\text{A.7})$$

Being an intersection of closed half-spaces, S° is always closed and convex.

Besides (A.1), another function associated with S is its *gauge*

$$\gamma_S(q) := \inf_{t>0, q \in tS} t$$

(knowing that $\gamma_S(q) = +\infty$ if $tq \in S$ for no $t > 0$). Polarity establishes a duality correspondence between σ and γ :

$$\text{If } S \text{ is closed and convex, then } \gamma_S = \sigma_{S^\circ} \text{ and } \sigma_S = \gamma_{S^\circ}$$

(note: if S is the unit-ball of some norm, (A.7) defines the dual norm, namely the one having S° as unit-ball).

When S is a cone, (i.e. $tx \in S$ whenever $x \in S$ and $t > 0$), S° is not changed if the righthand side “1” of (A.7) is replaced by any nonnegative number, say 0. Thus, the polar cone of a cone K is the closed convex cone

$$K^\circ := \{y : y^\top x \leq 0 \text{ for all } x \in K\} = \{y : \sigma_K(y) \leq 0\} \quad (\text{A.8})$$

of directions making an obtuse angle with all directions in K . Note: the polar of K is its normal cone at 0.

The polar of K° , i.e. the bipolar $K^{\circ\circ}$ of K , is just the closed convex hull of K (i.e. $K^{\circ\circ} = K$ if K is a closed convex cone). The gauge and support function of a closed convex cone are clearly either 0 or $+\infty$: indeed

$$\text{For a closed convex cone } K, [\gamma_{K^\circ}(y) =] \sigma_K(y) = \begin{cases} 0 & \text{if } y \in K^\circ \\ +\infty & \text{otherwise.} \end{cases}$$

When K is a polyhedral cone, namely $K = \{x \in \mathbb{R}^n : Ax \leq 0\}$, Farkas’ lemma states that $K^\circ = \{A^\top \lambda : \lambda \in \mathbb{R}_+^m\}$.

References

1. Balas, E.: Disjunctive programming. *Ann. Discrete Math.* **5**, 3–51 (1979)
2. Balas, E., Cornuéjols, G., Ceria, S.: A lift-and-project cutting plane algorithm for mixed 0-1 programs. *Math. Prog.* **58**, 295–324 (1993)
3. Balas, E., Perregaard, M.: Lift-and-project for mixed 0-1 programming: recent progress. *Discrete Appl. Math.* **123**, 129–154 (2002)
4. Bonami, P.: Etude et mise en œuvre d’approches polyédriques pour la résolution de programmes en nombres entiers ou mixtes généraux. PhD thesis, Université de Paris 6, 2003
5. Bonami, P.: Personal communication, 2004
6. Ceria, S., Soares, J.: Disjunctive cuts for mixed 0-1 programming: duality and lifting. Technical report, GSB, Columbia Univ., 1997
7. Ceria, S., Soares, J.: Convex programming for disjunctive convex optimization. *Math. Prog.* **86** (3), 595–614 (1999)
8. Cook, W., Kannan, R., Schrijver, A.: Chvátal closures for mixed integer programs. *Math. Prog.* **47**, 155–174 (1990)
9. Gomory, R.: An algorithm for the mixed integer problem. Technical Report RM–2597, The Rand Corporation, 1960
10. Hiriart-Urruty, J.-B., Lemaréchal, C.: *Fundamentals of convex analysis*. Springer Verlag, Heidelberg, 2001
11. Rey, P.A.: Análise convexa e métodos Lift-and-Project para programação inteira. PhD thesis, Dept. of Electrical Engineering, PUC, Rio, Brazil, 2001
12. Rockafellar, R.T.: *Convex analysis*. Princeton University Press, 1970